

# More precipitation may increase pollution of inland and coastal waters

October 20th, 2016



**In many parts of the world, water quality of inland streams, rivers and coastal waters suffers from pollutants from population and (agricultural) land use. This pollution may change because future precipitation will change, and so will pollutant transport to water bodies, and dilution of pollutants during high and low flows.**

**The impacts of climate change (precipitation and temperature) and land use on the daily microbial contamination of surface water were investigated for two catchments in the west of Ireland. This was done for an intermediate and a high-end scenario of climate change, and a potential worst-case land use scenario, for the period 2041 - 2060. Under the scenarios of climate change, changes in precipitation typically follow projected changes in future**

precipitation, with a stronger effect for the high-end than the intermediate scenario. A realistic worst-case land use scenario was used, based on the dominance of pasture and livestock in the study area: the cattle/sheep populations of 1998, when densities were at their highest level in the past 20 years. Manure applied to land as fertilizer and manure deposited from livestock grazing are major diffuse contaminant sources in the studied catchments. In addition, increase of effluent discharges from wastewater treatment plants and septic systems was assumed reflecting an estimated increase in human population of 17%.

The results show that daily microbial load will likely increase on an annual basis, following seasonal variations in precipitation and streamflow. Future winters represent periods of increased risk: increases in winter rainfall (intensity and frequency) according to the climate change scenarios produced significant effects on the transport of microbial pollutants. The land use scenario produced significant increases in microbial load compared with the current situation, across all seasons and flow conditions. The impacts are magnified when combined with the climate change scenarios.

It is generally expected that as climate changes, a greater proportion of annual precipitation will occur in larger magnitude events. More intense precipitation will contribute a greater fraction to direct runoff and may also cause a nonlinear increase in sediment erosion and pollutant loading. Within Ireland, the frequency of heavy precipitation events also shows notable increases, in particular for winter. Under the high-end scenario the frequency of very wet days (over 20 mm of precipitation) may increase by more than 20% in winter between now and halfway this century.

Future variations in land use/management may be as important as the effects of climate change on in-stream microbial pollutant loads. This highlights the need to develop adaptive land use/management strategies that are capable of reducing microbial source loading and to understand catchment characteristics that may be controllable in the future.

Source: Coffey et al., 2016. Regional Environmental Change 16: 2111 - 2128

Photo: Keith Ewing ([www.stckr.com](http://www.stckr.com))