

Future changes of wind energy potential over Europe

March 5th, 2016



By 2020, the European Commission aims to produce 15.7% of the EU's electricity from wind. A further increase of wind power capacity is expected for the upcoming decades. With respect to wind energy potential, it is more likely than not that this potential will increase over Northern and Central Europe on an annual basis. This is concluded from a study based on a large number of global climate models (GCMs). For Southern Europe, a likely decrease of mean annual wind energy potential is projected. These results agree with previous studies that indicate that future wind energy potential increases over Northern Europe, especially during winter, and decreases over Southern Europe.

For most of Europe (except for the Mediterranean area), wind energy potential is projected to increase for the winter and decrease for the summer. As a result, intra-annual variability of wind energy potential may strongly increase for most of Europe. In general, signals are stronger for 2061–2100 compared to 2021–2060 and for the scenario with strongest global warming (RCP8.5 compared to RCP4.5). A higher variability can potentially lead to a lower reliability of wind energy as an alternative energy source in future decades.

The differences in future projections of different models are large, however. Uncertainty about the magnitude of future changes is especially high for large parts of Germany, Scandinavia, and Eastern Europe. Uncertainties are much lower for Southwestern Europe: the strong decrease of future wind energy potential over Southwestern Europe is significant at the 95% confidence level.

The study was based on 22 global climate models (GCMs), and the model results were downscaled to the regional scale by a statistical method. In this study future projections were computed for two time periods (2021–2060 and 2061–2100); these projections were compared with the reference period 1961–2000. Two scenarios of climate change were considered: a scenario in which fossil CO₂ emissions are assumed to increase until 2040 and permanently decrease thereafter (the so-called RCP4.5 scenario), and a scenario in which the fossil CO₂ emissions strongly rise until 2100 (the RCP8.5 scenario).

Source: Meyers et al., 2016. *International Journal of Climatology* 36: 783–796.

Photo: Andreas Klink Johannsen (www.flickr.com)